

## Expression of *Cucumber mosaic cucumovirus* coat protein in *Escherichia coli* and production of specific polyclonal antiserum

Ali. M. El-Borollosy<sup>1</sup> and Sabah M. Hassan<sup>2</sup>

Department of Agricultural Microbiology<sup>1</sup> and Department of Genetics<sup>2</sup>, Faculty of Agriculture, Ain Shams University, P.O. Box 68, Hadayek Shubra 11241, Cairo, Egypt

Received: November 22 2013

Accepted: December 22 2013

### ABSTRACT

*Cucumber mosaic cucumovirus* (CMV) was isolated from naturally infected cucumber plants serologically depending on indirect enzyme-linked immunosorbant assay (I-ELISA) and biologically by mechanical inoculated on *Chenopodium amaranticolor* as a local lesions host and maintained on *Nicotianatabacum* cv. (White Burley). Infected tobacco plant tissue used to perform immunocapture reverse transcriptase polymerase chain reaction (IC-RT-PCR) to isolate the viral coat protein gene (*cp*). The agarose gel analysis of the PCR product indicated a single band with 657 bp length, which is the expected size for the CMV *cp*. The isolated CMV *cp* gene has been ligated into the PinPoint™ Xa-1 T-Vector depending on the T-A cloning principle. The new plasmids were transformed into competent *Escherichia coli* cells, further the gene integration success and orientation were screened by isolation of plasmids from transformed bacteria and treatment with *Bgl*III and *Bam*HI restriction enzymes. It was found that total plasmid length was 3331 bp without insert, while fragment separated after restriction digestion was approximately 660 bp. The PCR product of the minipreparation indicated a single band with 657bp which is the expected size for the CMV *cp* gene. After induction of CMV coat protein (CP) expression using Isopropyl β-D-1-thiogalactopyranoside (IPTG), transformed bacteria lysate analyzed using SDS-PAGE revealed a band with a molecular weight of 25KDa (the expected of CMV CP molecular weight). Western blotting analysis was carried out and confirmed that all the bands expected for viral CP react positively with CMV antiserum. The induced viral CP was purified from *E. coli* transformed cells using regenerated SoftLink™ Soft Release Avidin Resin yielding about 11 mg per 1 liter culture. Polyclonal antiserum produced in a New Zealand white rabbit by injecting of the produced fusion protein and Immunoglobulins G (IgGs) were purified from the collected antiserum. IgGs titration was performed using I-ELISA, the highest titers were 1:512 & 1:256 for antisera produced using fusion protein and purified virus, respectively. It was clearly observed that purified antiserum of *E. coli* viral fusion protein was more reactive and specific as it did not have any cross reactions with healthy tobacco plant sap.

**KEYWORDS:** *Cucumber mosaic cucumovirus*, Isolation, Coat protein gene, Cloning, Protein expression, Coat protein purification, Antiserum production, Immunoglobulins G purification, Evaluation.

### INTRODUCTION

*Cucumber Mosaic virus* (CMV, genus: *Cucumovirus*, family: *Bromoviridae*) is geographically wide spread and has been reported in Australia, North America, New Zealand, Europe and Africa (Roossinck 1999 and Sokhandan-Bashir *et al.*, 2012). It has four functional pieces of single stranded RNAs, packaged in icosahedral protein particles with molecular weight of 25 KDa, about 30 nm in diameter (Nouri *et al.*, 2012). The virus is transmitted by numerous species of aphid in a non-persistent manner (Brunt *et al.*, 1996 and Gildow *et al.*, 2008). The virus has an extremely wide host range and it induces severe symptoms. Douine *et al.* (1979) listed 775 plant species representing 365 genera and 85 families that are susceptible to CMV infection. In Egypt CMV infects cucurbit and solanaceous crops, banana and sugar beet causing remarkable yield loss (El-Affi *et al.*, 2007).

Serological methods especially the enzyme linked immunosorbant assay (ELISA) are widely used in the detection of viral infections (Portsmann and Kiessig, 1992). They are relatively sensitive, inexpensive, simple and suitable for the testing of many samples simultaneously (Clark, 1981). Additionally, more specific antibodies are a prerequisite for the application of the extremely sensitive technique immunocapture reverse transcriptase polymerase chain reaction (IC-RT-PCR) (Wetzel *et al.*, 1992; Komorowska and Malinowski, 2009).

\*Corresponding Author: Ali. M. El-Borollosy, Department of Agricultural Microbiology, Faculty of Agriculture, Ain Shams University, P.O. Box 68, Hadayek Shubra 11241, Cairo, Egypt

Generally, molecular methods such as RT-PCR are not suitable as routine tests for indexing large numbers of samples due to costs and the relative complexity of execution. As a consequence, serology has traditionally been the most used method of plant virus diagnosis in a large number of samples, using ELISA as the method of choice (Zimmermann *et al.*, 1990; Ling *et al.*, 2000; Fajardo *et al.*, 2007).

Recently, advances in recombinant DNA technology, coupled with its ease to manipulate and its rapid growing rate in a less expensive media had established *E. coli* as a leading host organism to produce high protein quantities of scientific interest (Ashoub *et al.*, 2009). Many of the expression systems have advantages and limitations but *E. coli* as a prokaryotic system was used widely because of its high expression levels of many heterologous proteins, low cost, efficient generation time, and fast high density cultivation (Sokhandan-Bashir *et al.*, 2012).

Virus purification is usually a labor-intensive procedure with varying, occasionally unsatisfactory results concerning purity and concentration of the final preparation (Ling *et al.*, 2007). The production of high quality virus-specific antiserum suitable for large scale of virus detection and based on virus purification procedures faces substantial drawbacks, *i.e.*, including complex virus infections, low yields of virus particles, contamination of antigens with plant proteins, presence of inhibitory compounds (Radaelli *et al.*, 2008; Khan *et al.*, 2012). On the other hand, it was found that the production of virus-specific antisera using recombinant proteins from cloned virus genes expressed in *E. coli* "which can overcome such difficulties" has been applied to produce a number of polyclonal antisera against coat protein (CP) of several plant viruses, *i.e.*, *Bean yellow mosaic potyvirus* (Hammond and Hammond, 1989), *Plum pox potyvirus* (Mattanovitch *et al.*, 1989), *Soybean mosaic potyvirus* (Liu *et al.*, 1993), *Cucumber mosaic cucumovirus* (Khan *et al.*, 2011; Sokhandan-Bashir *et al.*, 2012).

The aim of this study is to express CMV coat protein in *E. coli*, which was then used for the production of specific polyclonal antibodies. As antibodies with higher specificity are a crucial reagents in carrying out serological tests for screening against viruses in plant material such as seeds and other propagation material and also for field plants testing.

## MATERIAL AND METHODS

### Virus isolation:

Fifty samples of naturally infected cucumber plants (Collected from the open field of Faculty of Agriculture, Ain Shams University, Cairo, Egypt) showing mosaic, mottling and malformation were used for virus isolation. According to Koenig (1981) I-ELISA was performed on samples using polyclonal antibodies specific for CMV (Agdia Inc., USA). Samples giving CMV positive results were mechanically inoculated on *Chenopodium amaranticolor*. Local lesions produced were used as a source of single lesion isolation, and finally the virus was maintained in *Nicotiana tabacum* cv. White Burley under greenhouse conditions (28°C ±2).

### Isolation CMVcp gene using IC-RT-PCR:

Immunocapturing, the isolation of CMV RNA and cDNA synthesis using IC-RT-PCR was performed using infected tobacco leaf collected 15 days post CMV inoculation according to Minafera and Hadidi (1994) as follows: the wells of an ELISA plate were coated by adding 200 µl of IgGs (diluted to 10-1 using coating carbonate buffer (pH 9.6)) and incubated 4 hrs at 37°C, after that, plate was washed 3 times with PBST, 5 min each. Plant tissues were ground in sample buffer (1 gm : 5 ml), then centrifuged at 3000 rpm/10 min at 4°C, 200 µl of the supernatant were added to each pre-coated wells and incubated overnight at 4°C, then plate was washed 3 times, 5 min each with PBST.

Twenty five of pre-heated transfer buffer (10 mM Tris-HCl (pH 8.0) containing 1% Triton X-100) were added to each well and kept at 65°C for 5 min. Aliquots of 5 µl from the resulting released RNA solutions were immediately transferred to microcentrifuge tube. To each tube the following components were added and incubated at 42°C/2 h for cDNA synthesis: 1 µg of reverse coat protein gene primer, 4 µl of 5X first strand buffer (250 mM Tris-HCl (pH 8.3), 375 mM KCl, 15 mM MgCl<sub>2</sub>), 3 µl of 0.1M dithiothreitol (DTT), 5 µl of 0.3M 2-ME, 2.5 µl of 10 mM of each deoxynucleotide triphosphate (dNTPs), 1 µl RNasin (40 unit) and 1 µl of Molny Murine Leukemia Virus (M-MLV) (200 unit). Final product was used immediately for PCR reaction.

The following primers (Invitrogen Corp., USA) designed depending on CMV *cp* gene sequences collected from GenBank web site (<http://www.ncbi.nlm.nih.gov>) were used for the amplification of CMV *cp* gene were:

5'ATGGACAAATCTGAATCAAC3' (Sense) 5'TCAAACCTGGGAGCACCCAG3' (Antisense).

PCR was performed as described by **Ghosh et al. (2002)** using PerkinElmer Cetus Thermal Cycler PE 9700 (PerkinElmer Inc., USA). Five  $\mu\text{l}$  from resulting cDNA were transferred to tube containing 45  $\mu\text{l}$  PCR reaction mixture. PCR program was 94°C initial melting for 3 min followed by 35 cycles of 94°C/1 min, 55°C/1 min, 72°C/2 min and 72°C/10 min final extension.

For PCR product analysis, 1.5% agarose gel was used and electrophoresis was carried out in Sub-Cell DNA apparatus (Bio-Rad® Lab., USA) at 80 V. The amplified gene band was visualized on an UV Transilluminator and photographed by Gel Documentation System (AlphaImager® TM1220, Documentation and Analysis system, Canada).

#### **Cloning of CMV *cp* gene:**

The virus *cp* gene (PCR product) was ligated with the PinPoint™ Xa-1 T-Vector (Promega Corp., USA) depending on the T-A cloning principle (**Holton and Graham, 1991**) and according to manufacturer instructions manual. The gene was cloned with the vector in-frame with the fragment of biotin-binding protein (BBP) as mentioned by **Abou Zeid et al. (2002)**.

The recombinant plasmids were transferred into competent *E. coli* strain BL21 cells and successfully transformed bacteria were selected depending on blue/white colonies screening. Transformation and selection were carried out as described by **Hanahan (1985)**.

#### **Minipreparation and restriction enzyme digestion:**

The recombinant plasmid was isolated from *E. coli* transformed cells according the miniprep procedure described by **Sambrook et al. (1989)**. The preparation was subjected to restriction analysis to confirm *cp* gene orientation prior to protein expression. As described by **Sambrook et al. (1989)** with some modifications, 15  $\mu\text{l}$  of the recombinant plasmid was mixed with 2  $\mu\text{l}$  from both *Bgl*III and *Bam*HI restriction enzymes (Promega Corp., USA), the total volume was raised to 30  $\mu\text{l}$  by deionized water. The reaction was incubated at 37°C for 2 h, and then 20  $\mu\text{l}$  of the digested product was mixed with 6  $\mu\text{l}$  agarose gel loading buffer and electrophorised on 1% agarose gel.

PinPoint™ Xa-1 original plasmids (without the CMV *cp* gene) and recombinant plasmids with the CMV *cp* gene in addition to PCR product of the minipreparation using viral *cp* gene specific primers were loaded to the same gel as controls.

#### **Induction of *cp* gene expression and detection of the expressed protein:**

As recommended by manufacture (Promega) instructions, culture of *E. coli* carrying the PinPoint™ Xa-1 expression fusion was started by inoculating 5 ml of LB medium containing 100  $\mu\text{g/ml}$  ampicillin with a 24 h colony and culture was incubated overnight at 37 °C on shaker. Overnight culture was diluted to 1:100 by adding 50  $\mu\text{l}$  of culture to 5 ml of LB medium containing 2  $\mu\text{M}$  biotin, 100  $\mu\text{g/ml}$  ampicillin and incubated for 1 h at 37 °C on shaker. For protein expression induction, 100  $\mu\text{M}$  isopropyl  $\beta$ -D-thiogalacto-pyranoside (IPTG) was added to culture, incubated for 4 h at 37 °C with shaking. Bacterial cells were then collected by centrifugation and further subjected to subsequent freezing/thawing, sonication and centrifugation treatments for protein extraction (**Sambrook et al., 1989**). The proteins were then separated by 12 % SDS polyacrylamide gel electrophoresis (SDS-PAGE) (**Laemmli, 1970**), electro-blotted to nitrocellulose membranes and assayed by Western blotting immunoassay using polyclonal CMV coat protein specific antibodies (**Iracheta-Cardenas et al., 2002**).

#### **Protein purification:**

For the purification of large protein quantities, one liter of transformed bacteria culture in LB broth containing 2  $\mu\text{M}$  biotin and 100  $\mu\text{g/ml}$  ampicillin was prepared. The protein expression was induced by adding 100  $\mu\text{M}$  IPTG to the bacterial culture and incubated 4 h at 37 °C on shaker. The cells were disrupted by sonication according to **Sambrook et al. (1989)** and centrifuged at 8000 rpm for 10 min at 4 °C and the fusion protein was then purified using 3 ml regenerated Soft Link™ Soft Release Avidin Resin according to Promega instructions. The eluted fusion protein was quantified by the Bradford assay (**Bradford, 1976**) at  $A_{595}$  nm using spectrophotometer. The protein was then analyzed and separated by electrophoresis in 12 % SDS-PAGE and detected using Western blotting immunoassay. As a control a purified CMV preparation was also loaded to gel.

#### **Antisera production and evaluation:**

Depending on the method described by **Madhubala *et al.* (2005)** polyclonal antiserum produced in a New Zealand white rabbit by injecting of the produced fusion protein and a purified CMV preparation as a control (purified as described by **Lot *et al.*, 1972**) (1 mg of virus with Freund's incomplete adjuvant, 1 : 1) intramuscularly six times at 10 days interval. The animal was bled 15 days after the last injection and the antiserum was collected.

Immunoglobulins G (IgGs) purification was carried out as mentioned by **Clark and Adams (1977)**. Final dialyzed proteins were loaded on 10 ml of diethylaminoethyl (DEAE) cellulose column (Whatman Inc., USA), eluted using half strength phosphate buffer saline (PBS) and collected in microtube. Absorption of fractions was measured at 280 nm and concentration of IgGs was adjusted to 1 mg/ml ( $OD_{280}=1.46$ ) using half strength PBS, then stored at 4°C. IgGs dilution end points were determined using I-ELISA against clarified infected tobacco sap and healthy tobacco sap was used as a control.


## RESULTS

### Isolation of CMV virus:

CMV was isolated from naturally infected cucumber plants showing mosaic, mottling and malformation. Samples gave positive I-ELISA results ranging from 0.897 to 1.095 were mechanically inoculated on *Ch.Amaranticolor* leaves and gave chlorotic local lesions. Lesions were extracted and inoculated on *N.tabacum* cv. White Burley for maintenance.

### Isolation of CMV *cp* gene using IC-RT-PCR technique:


Infected tobacco plant tissues were used to perform IC-RT-PCR to isolate the viral *cp* gene. The agarose gel electrophoresis of the PCR product indicated a single band sized 657bp, which is the expected size for the CMV *cp* gene (**Figure 1**).



**Figure 1.** IC-RT-PCR results for the isolation of CMV *cp* gene (Lane 1-2). M: DNA Ladder (Promega, USA).

### Cloning of CMV *cp* gene:

The PCR isolated CMV *cp* gene has been ligated into the PinPoint™ Xa-1 T-Vector depending on the T-A cloning principle. The new recombinant plasmid vector was transformed into competent *E. coli* cells, further the gene integration success and orientation were proved by isolating the recombinant PinPoint™ Xa-1 T-Vector from transformed bacteria and treating it with *Bgl*II and *Bam*HI restriction enzymes, which restricted to certain sites through the polylinker region of the plasmid vector. It was found that total plasmid length was 3331 bp without the insert, while the fragment separated after restriction digestion was approximately 660pb in length (gene length plus some added nucleotides after the enzymes cutting sites) compared to the PCR product using the recombinant PinPoint™ Xa-1 T-Vector as a template, which gives a single band with 657bp length, the expected size for the CMV *cp* gene (**Figure 2**).




**Figure 2.** Restriction analysis to check for the right integration of the *cp* gene into PinPoint™ Xa-1 T-Vector: The empty PinPoint™ Xa-1 plasmid vector (Lane 1), the recombinant PinPoint™ Xa-1 T-Vector (containing CMV *cp* gene)(Lane 2), the recombinant PinPoint™ Xa-1 T-Vector cut with *Bgl*II and *Bam*HI restriction enzymes (Lane 3 & 4) and PCR product for CMV *cp* gene using the recombinant PinPoint™ Xa-1 T-Vector as a template (Lane 5). M: DNA Ladder (Promega, USA).

**Expression, detection and purification of the CMV coatprotein transformed *E. coli* cells:**

Transformed bacteria lysate analyzed using SDS-PAGE revealed a band with a molecular weight of 25KDa(Figure 3A)after induction of protein expression using IPTG.

To produce large amount of the fusion protein, large scale culture from transformed *E. coli* was prepared and protein was purified depending on affinity purification using SoftLink™ Soft Release Avidin Resin. About 11 mg of expressed protein was purified from 1 liter of bacterial culture. The purified protein gave a band with molecular weight of 25KDa, which is expected to CMV coat protein (Figure 3A).

Result was confirmed by transferring gel bands to nitrocellulose membrane for Western blotting analysis, and all the bands expected for viral CP (25 KDa) react strongly with CMV specific antiserum (Figure 4B).



**Figure 3.** (A) SDS-PAGE analysis for the lysate of *E. coli* cells: Transformed *E. coli* lysate (Lane 1), untransformed *E. coli* lysate as a control (Lane 2), purified fusion protein(Lane 3) M: Protein marker (Promega, USA). (B) Western blotting analysis showing positive reaction with the viral protein bands obtained from the transformed *E. coli* lysate.

**Antisera production and evaluation:**

IgGs titration results were illustrated in **Table (1)**, the highest titers were 1:512 & 1:256 for antisera produced using fusion protein or purified virus preparation, respectively. It was clearly observed that purified antiserum of viral fusion protein produced from bacterial cells was more reactive as it gave higher I-ELISA values and more specific for not giving any cross reactions with healthy tobacco plant sap.

**Table 1. Titration and evaluation of CMV IgGs produced from fusion protein and purified virus preparation against infected and healthy tobacco sap**

Dilution	I-ELISA values at 405 nm*							
	IgGs <sub>1</sub>			IgGs <sub>2</sub>				
	I	R	H	R	I	R	H	R
Crude	0.914	+	0.122	-	0.820	+	0.404	+
1:1	0.825	+	0.100	-	0.801	+	0.311	-
1:2	0.810	+	0.099	-	0.766	+	0.209	-
1:4	0.805	+	0.091	-	0.677	+	0.259	-
1:8	0.787	+	0.088	-	0.660	+	0.199	-
1:16	0.731	+	0.069	-	0.611	+	0.091	-
1:32	0.711	+	0.065	-	0.589	+	0.100	-
1:64	0.677	+	0.062	-	0.579	+	0.095	-
1:128	0.649	+	0.059	-	0.490	+	0.086	-
1:256	0.550	+	0.060	-	0.441	+	0.084	-
1:512	0.504	+	0.059	-	0.064	-	0.069	-
1:1024	0.200	-	0.049	-	0.061	-	0.065	-
Purified**	1	1.353		+	1.077		+	
	2	1.042		+	0.924		+	

\* Each ELISA result (R) was the average of three readings.

\*\* Reaction of CP fusion protein (1) and crude purified virus (2) with crude antiserum.

Note: Reaction of infected (I) and healthy (H) tobacco leaves sap with antisera produced by rabbit immunization with CP fusion protein (IgGs<sub>1</sub>) and from purified virus preparation (IgGs<sub>2</sub>).

## DISCUSSION

*Cucumber mosaic cucumovirus* (CMV), one of the most common virus infecting some economical plants in Egypt, the virus infects important crops like sugar beet, banana and cucurbits causing significant loss in yield quality and quantity (El-Affiet *et al.*, 2007; El-Borollosy and Oraby, 2012).

Such virus is difficult to diagnose due to the periodic changes in symptoms and also for the need of high quantity of antiserum with higher specificity to perform routine serological tests for indexing large numbers of samples in field (Sokhandan-Bashir *et al.*, 2012).

The CMV *cp* was isolated depending on IC-RT-PCR technique and then this gene has been ligated into the PinPoint™ Xa-1 T-Vector depending on the T-A cloning principle. The obtained recombinant PinPoint™ Xa-1 T-Vector was transformed into competent *E. coli* cells, further the *cp* gene integration was proved by restriction enzymes analysis. The PCR product of the isolated recombinant PinPoint™ Xa-1 T-Vector from the transformed *E. coli* cells produced a single band with 657bp, which is the expected size for the CMV *cp* gene. There are numerous reports of the expression of plant viruses CP in *E. coli* (Hammond and Hammond, 1989; Mattanovicht *et al.*, 1989; Liu *et al.*, 1993; Khan *et al.*, 2012; Sokhandan-Bashir *et al.*, 2012).

Transformed bacteria lysate analyzed using SDS-PAGE revealed a band of CMV CP with a molecular weight of 25KDa. Western blotting analysis was carried out and confirmed that all the bands expected for viral CP react positively with CMV antiserum. Such work agreed with that performed by Khan *et al.* (2012) and Sokhandan-Bashir *et al.* (2012).

The induced viral CP was purified from *E. coli* transformed cells using regenerated SoftLink™ Soft Release Avidin Resin yielding About 11 mg of protein from 1 liter of bacterial culture. This result is a promising result when compared with that obtained by Khan *et al.* (2012) who produced a total yield of about 8 mg CMV CP per 1 liter of cell culture.

Polyclonal antiserum produced by rabbit immunization and IgGs were purified from the collected antiserum. Depending on I-ELISA it was clearly observed that purified antiserum of *E. coli* viral fusion protein has higher dilution end point, having more reactivity for giving higher I-ELISA values, and

specificity as it did not give any cross reactions with healthy tobacco plant sap which can improve the performance of the antiserum.

**Barbieriet al. (2004)** and **Sokhandan-Bashir et al. (2012)** concluded, based on observations with *Watermelon mosaic potyvirus* and *Cucumber mosaic cucumovirus*, respectively that antibodies produced against expressed CPs in *E. coli* tend to be more specific, reducing the occurrence of unexpected heterologous reactions.

The use of fusion CPs as an immunogen to develop antibodies has been of great value particularly for viruses that are difficult to purify (**Petrovicet al., 2003**) and for those that are subject to degradation during the purification process (**Barbieriet al., 2003**).

## REFERENCES

- AbouZeid, A.A., A. Barakat and H.A. Amin (2002)**. Expression and Purification of the Coat Protein Gene (P25) of *Citrus tristeza virus* in *E. coli*. **Al-Azhar J. Microbiology** **58**: 1-9.
- Ashoub, A., Mervat M.M. El-Far and M.I. Salama (2009)**. Expression of Sweet potato feathery mottle virus coat protein and production of specific antiserum for diagnostic aspects. **Egypt. J. Genet. Cytol.** **38**:119-126.
- Barbieri, M.R., M.G. Carvalho, E.M. Zambolim and F.M. Zerbini (2003)**. Expressaoem *Echerichia coli* da proteinacapsidal do watermelon mosaic virus e producao de anti-soro. **Fitopatol Bras** **29**:215-219.
- Bradford, M.N. (1976)**. A rapid and sensitive method for quantification of microgram quantities of protein utilizing the principle of protein-dye binding. **Anal Biochem.** **72**: 248–254.
- Brunt, A., K. Crabtree; M. Dallwitz, A. Gibbs and L.Watson (1996)**. Viruses of Plants. Description and Lists from the VIDE Database, pp. 477-483. **CAB International Wallingford, UK**.
- Clark, M.F. (1981)**. Immunosorbent assays in plant pathology. **Annual Review of Phytopathology** **19**: 83-106.
- Clark, M.F. and N.E. Adams (1977)**. Characteristics of the microtiter plate method of enzyme-linked immunoassay (ELISA), for the detection of plant viruses. **J. Gen. Virol.** **37**: 475-483.
- Douine, L., J.B. Quiot, G. Marchoux and P. Archange (1979)**. Recensement des especes vegetales sensible au virus de la mosaïque du concombre. Etude bibliographique. **Annales de Phytopathologie** **11**: 439-475.
- El-Affifi, Sohair I., A.M. El-Borollosy and S.Y.M. Mahmoud (2007)**. Tobacco callus culture as a propagating medium for *Cucumber mosaic cucumovirus*. *Inter. J. Virology.* 3: 73-79.
- El-Borollosy, A.M. and Mona M. Oraby (2012)**. Induced systemic resistance against *Cucumber mosaic cucumovirus* and promotion of cucumber growth by some plant growth-promoting rhizobacteria. **Ann. of Agric. Sci.** **57**: 91-97.
- Fajardo, T.V.M., D.R. Barros, O. Nickel, G.B. Kuhn and F.M. Zerbini(2007)**. Expression of Grapevine leafroll-associated virus 3 coat protein gene in *Escherichia coli* and production of polyclonal antibodies. **Fitopatologia Brasileira** **32**: 496-500.
- Ghosh, S.B., L.H.S. Nagi, T.R. Ganapathi, S.M. Paul Khurana and V.A. Bapat (2002)**. Cloning and sequencing of *Potato virus Y* coat protein gene from an Indian isolate and development of transgenic tobacco for PVY resistance. *Current Science* 82: 7-10.
- Gildow, F.E., D.A. Shah and W.M. Sackett (2008)**. Transmission efficiency of Cucumber mosaic virus by aphids associated with virus epidemics in snap bean. **Phytopathology** **98**: 1233-1241.
- Hanahan, D. (1985)**. Techniques for transformation of *E. coli*. In DNA cloning: A Practical Approach(ed. D.M. Glover), vol. 1 pp. 109-135. **IRL Press, Oxford, UK**.
- Hammond, J. and R.W. Hammond (1989)**. Molecular cloning, sequencing and expression in *Escherichia coli* of the Bean Yellow Mosaic Virus coat protein gene. **J. Gen. Virol.** **70**: 1961-1974.
- Holton, T.A. and M.W. Graham (1991)**. A simple and efficient method for direct cloning of PCR products using ddT-tailed vectors. **Nucleic Acids Res.** **19**: 1156.
- Iracheta-Cardenas, M.M., I.H. Almeyda-Leon and K.L. Manjunath (2002)**. Molecular cloning and expression of the p25 gene of two Mexican isolates of *Citrus tristeza virus*. **Rev Mex Fitopatol****20**:174-181.
- Khan, S., A.T. Jan, B. Mandal and Q.M.R. Haq (2012)**. Immunodiagnosics of cucumber mosaic virus using antisera developed against recombinant coat protein. **Archives of Phytopathology and Plant Protection** **45**: 561-569.

- Koenig, R.C. (1981).** Indirect ELISA methods for broad specificity detection of plant viruses. *J. Gen. Virol.* 55: 53-62.
- Komorowska, B. and T. Malinowski (2009).** The attempts to produce antiserum against Apple stem pitting virus coat protein (ASPV CP) obtained in prokaryotic and eukaryotic expression systems. *Journal of Fruit and Ornamental Plant Research* 17: 21-30.
- Laemmli, U.K. (1970).** Cleavage of structural proteins during the assembly of the head of bacteriophage T<sub>4</sub>. *Nature* 227: 280-285.
- Ling, K.S., H.Y. Zhu, Z.Y. Jiang and D. Gonsalves (2000).** Effective application of DAS-ELISA for detection of grapevine leafroll associated closterovirus-3 using a polyclonal antiserum developed from recombinant coat protein. *European Journal of Plant Pathology* 106: 301-309.
- Liu, J., X. Peng, L. Li and K. Mang (1993).** Cloning of coat protein of *Soybean mosaic virus* and its expression in *Escherichia coli*. *Chin. J. Biotechnol.* 9: 143-149.
- Lot, H., G. Jonard and K.E. Richards (1977).** Cucumber mosaic virus RNA 5. Partial characterization and evidence for no large sequence homologies with genomic RNAs. *FEBS Letters* 80: 395-400.
- Madhubala, R., V. Bhadramurthy, A.I. Bhat, P.S. Hareesh, S.T. Rethesh and R.S. Bhai (2005).** Occurrence of *Cucumber mosaic virus* on vanilla (*Vanilla planifolia* Andrews) in India. *J. Biosci.* 30: 339-350.
- Mattanovich, D., G. Himmler, M. Laimer, E. Maiss, F. Regner, A. Da Camara Machado, V. Hanzer, R. Casper and H.W. Katinger (1989).** Expression of the *Plum pox virus* coat protein region in *Escherichia coli*. *Virus Genes* 2: 119-127.
- Minafera, A. and A. Hadidi (1994).** Sensitive detection of *Grapevine virus A, B* or leafroll associated III from viruliferous mealy bugs and infected tissues by cDNA amplification. *J. Virol. Methods* 47:175-188.
- Nouri, S., B.W. Falk and R.L. Groves (2012).** A new satellite RNA is associated with natural infections of cucumber mosaic virus in succulent snap bean. *Arch. Virol.* 157: 375-377.
- Petrovic, N., B. Meng, M. Ravnkar, I. Mavric and D. Gonsalves (2003).** First detection of Rupestris stem pitting associated virus particles by antibody to recombinant coat protein. *Plant Disease* 87:510-514.
- Porstmann, T. and S.T. Kiessig (1992).** Enzyme immunoassays techniques. An overview. *Journal of Immunological Methods* 150: 5-21.
- Radaelli, P., T.V.M. Fajardo, O. Nickel, M. Eiras and G. Pio-Ribeiro (2008).** Production of polyclonal antisera using recombinant coat proteins of *Grapevine leafroll-associated virus 2* and *Grapevine virus B*. *Pesq. agropec. bras.* 43: 1405-1411.
- Roossinck, M.J. (1999).** Cucumoviruses (Bromoviridae)-general features, In: "Encyclopedia of Virology, 2nd edition", pp. 315-320, (Granoof L. & Webster R.G. eds). Academic Press, San Diego, USA.
- Sambrook, J., E.F. Fritsch and T. Maniatis (1989).** Molecular Cloning. Cold Spring Harbor Lab. Press, USA.
- Sokhandan-Bashir, N., A. Rostami and D. Koolivand (2012).** Towards Preparation of Recombinant Antibody to *Cucumber Mosaic Virus*: Expression of the Coat Protein in *Escherichia Coli*. 2<sup>nd</sup> Inter. Conf. on Advances in Biotech. and Pharmaceutical Sci. June 30 July 1, 2012 Bali, Indonesia.
- Wetzel, T., T. Candresse, G. Macquaire, M. Ravelonandro and J. Dunez (1992).** A highly sensitive immunocapture polymerase chain reaction method for plum pox potyvirus detection. *Journal of Virological Methods* 39: 27-37.
- Zimmermann, D., P. Bass, R. Legin and B. Walter (1990).** Characterization and serological detection of four closterovirus-like particles associated with leafroll disease on grapevine. *Journal of Phytopathology* 130: 205-218.