

Physicochemical Composition and Sanitary Quality of Pasteurized Milk Marketed in Western Algeria

Habiba Fernane¹, Aicha Tir Touil², Hama Benbarek² and Mokhtar Benchohra^{1*}

¹Department of Biomedicine, Institute of Veterinary Science, Ibn Khaldoun University, Tiaret 14000, Algeria

²Department of Biology Science, Faculty of Natural and Life Sciences, Mustapha Stambouli University, Mascara 29000, Algeria

Received: July 28, 2016

Accepted: October 10, 2016

ABSTRACT

This study was carried out to assess the safety and to check for thermo-resistant bacteria of pasteurized cow milk. Physicochemical and bacteriological tests were performed on 100 pasteurized milk samples. The acidity, density, fat, total dry extract (TDE), and the defatted dry extract (DDE) were studied; also, germs were researched according to the national standards. Mesophilic flora count was high; it varied between 10^4 and 1.4×10^6 , the total coliforms (TC) and fecal coliforms (FC) content was elevated. 61% of sampled milk was contaminated with *Staphylococcus aureus*, 34% by Sulphite-reducing *Clostridia* (SRC) and 100% by fecal *Streptococci* (FS). The overall results obtained suggested that pasteurized milks were of poor hygienic quality and were not conformed to the Algerian standard; in infringement of the current sanitary regulation. Thus, there is urgent need to test and monitor the milk chain periodically, after pasteurization and packaging.

KEYWORDS: Milk hygienic Quality, Milk composition, Pasteurization, Contamination, Thermo-Resistant Bacteria

INTRODUCTION

Milk is an important component of our daily diet; it is an important source of minerals, carbohydrate, protein, lipid and vitamins. Due to the growing human needs, milk technology has pioneered methods to better preserve the milk for a long time, such pasteurization and sterilization.

However, milk is a nutritious medium that can support growth of a large selection of bacterial contaminants; bacteria are able to use proteins, fats, carbohydrates and vitamins in milk for their growth and metabolism [1]. Hence, milk should not be consumed or used in dairy products without a former pasteurization.

Pasteurization process of milk is a heat treatment intended to reduce the number of any harmful microorganisms to a level at which they do not constitute a significant health hazard; reduce the level of undesirable enzymes and spoilage bacteria, and thus increase the keeping quality; achieve the preceding two goals while maintaining the nutritional integrity of the original product [2].

Nevertheless, some agents responsible for zoonosis can be transmitted to the human by milk consumption even if pasteurized [3] so, control of pasteurized milk hygiene proves to be very important. As well, it may be taking into account the conditions in which the milk is produced, including the respect of hygiene mastery of manipulation, to avoid post pasteurization contamination.

In Algeria, a few works were focused on milk quality evaluation [4,5]. The present investigation aimed to spot light on the quality of pasteurized cow milk marketed in Tiaret area (Algeria); determined by physicochemical and bacteriological tests.

MATERIAL AND METHODS

Sources of milk samples

Milk samples were collected from GIPLAIT (public milk factory) of Tiaret District in Western Algeria. The study was conducted from December 2013 to May 2014, on 100 samples taken from 100 packets of pasteurized milk. Samples were stored at 6-8 °C till laboratory tests carried out after 4 hours.


Experimental protocol

Physicochemical analysis: Acidity was measured by titration with a solution of sodium hydroxide (N/ 9) in the presence of the phenolphthalein at 1 percent, as indicator [6]. Density was calculated by a thermolactodensimeter type Dornic. It is the ratio of masses of a same volume of milk and water at 20°C [7]. Fat was carried out by the butyrometer according to the method described by Abiazar [8]. Total dry extract (TDE) and the defatted dry

*Corresponding author: Mokhtar Benchohra, Department of Biomedicine, Institute of Veterinary Science, Ibn Khaldoun Univ Tiaret, Poste de l'Université BP 1, 14010 Tiaret, Algeria, E-mail:mok.benchohra@yahoo.fr

extract (DDE) expressed as a percentage mass, consist on evaporation method; residue was weighed thereafter [9].

Microbiological analysis: Germs were examined by total aerobic mesophilic flora (TAMF) at 30°C; enumeration was carried out on agar PCA (Pasteur Institute Algeria) [7].VRBL agar (Difco) was used to detect TC (Total coliforms) at 37°C and FC (Fecal coliforms) at 44°C [10]. Research of *S. aureus* was made according to Giraud [11] and count using Baird-parker agar (Difco) [12]. Determination of SRC (Sulphite-reducing *Clostridia*) was carried out according to Catsaras and Bourgeois [13]. Whereas, VF agar was used for detecting FS (Fecal streptococci) and counting was done on Roth and Litsky medium [14]. Conducted tests are summarized in the following flowchart:



RESULTS AND DISCUSSION

Physicochemical characteristics of milks

The results of analyses are summarized in table 1; they were discussed following the national standard [15].

Table 1: Physicochemical parameters in checked samples of pasteurized milk.

Parameters							Total No. of samples	Mean	Standard [15]
TA (°D)	17°	18°	20°	--	--	--	n=100	17.8°	16° - 19°
	(n=40)	(n=50)	(n=10)	--	--	--			
Density	1.029	1.030	1.031	1.031	1.031	1.031	n=100	1.030	1.027 - 1.035
	(n=10)	(n=20)	(n=20)	(n=30)	(n=10)	(n=10)			

Fat g/l	24 (n=20)	27 (n=30)	28 (n=30)	30 (n=20)	-- --	-- --	-- --	n=100	27.3	30 - 45
TDE g/l	109.5 (n=10)	115 (n=10)	116.5 (n=10)	118 (n=20)	118.5 (n=30)	119.5 (n=10)	120 (n=10)	n=100	117.2	125 - 130
DDE g/l	89.5 (n=20)	90 (n=10)	90.5 (n=20)	91 (n=10)	91.5 (n=10)	92 (n=20)	109.5 (n=10)	n=100	92.5	90 - 95

TA= Titratable acidity; °= Dornic degree; TDE= Total dry extract; DDE= Defatted dry extract;
n = number of samples

The large proportion of checking milk (90%) had a titratable acidity (TA) close to standard [15] except for 10 samples; showing a slight elevation. Acidity increase might be due to a high microorganism's number of milk and/or to microorganism's proliferation; that metabolism caused a high production of lactic acid, on one hand, and hygienic conditions during milking, on other [7]. According to Schmidt *et al.* [16] as raw milk increased in age TA increased to the upper level; also, augmentation of milk protein content has the same effect. Concerning milk density, it was in compliance with the national standard. About the fat content of milks, 80% of samples results ranged between 24 and 28g/l; which are below the reference value (Table 1). This is the most variable component of milk [17], its less level might be due to an excessive extraction of fat from raw milk in processing for producing butter. TDE rate of all results are substandard; the dray matter represents all the components of milk except for water and dissolved grasses. The less level of TDE can be explained by milk dilution. For the DDE, according to Veisseyre [18] one liter of milk contains 90 to 95 g; 90% of our results accord with these values which are within the regulatory limit, except ten samples; with high level (109.5g/l). This increase might be due to environmental and dietary factors.

Total mesophilic flora and hygienic quality

The results of bacteriological analyses are presented in table 2; they were discussed following the regulatory limit [12].

Table 2: Microbiological parameters in checked samples of pasteurized milk.

Total samples (N=100)	TAMF	Total coliforms	Fecal coliforms	Staphylococcus aureus	Sulphite-reducing Clostridia	Fecal Streptococci
	Number of colonies (CFU/ml)					
12	1.4×10 ⁶	10 ⁵	1.4×10 ⁴	Present	Present	150
12	1.2×10 ⁵	2.1×10 ⁴	1.6×10 ³	Present	Present	120
10	1.5×10 ⁵	10 ⁴	Absent	Present	Present	120
12	2.1×10 ⁵	1.8×10 ⁴	Absent	Absent	Absent	210
08	1.6×10 ⁴	3.1×10 ³	4.1×10 ²	Absent	Absent	1100
09	1.9×10 ⁵	1.8×10 ⁴	10 ³	Present	Absent	460
09	1.7×10 ⁴	200	Absent	Present	Absent	1100
09	2.6×10 ⁴	1.6×10 ²	Absent	Present	Absent	240
13	1.3×10 ⁴	1.3×10 ²	Absent	Absent	Absent	1100
06	10 ⁴	Absent	Absent	Absent	Absent	460
Standards	10 ⁵	10 ³	10 ²	Absent	Absent	Absent/0.1ml

Thus, the microbiological quality control of pasteurized cow milk samples is determined by the count of mesophilic aerobic bacteria, indicators who provide an assessment on the overall level of milk contamination [14]. TAMF permit to appreciating microbial pollution and general quality of food product [18, 19]. The fact that TAMF charges were highest, in 55% of samples, proved that milk pasteurization is not efficient. In practice, an efficient pasteurization process should kill most of pathogen bacteria; however, some bacteria can survive pasteurization and are called thermodurics [20]. Or a possible contamination of the product has been occurred during packaging processes. Also, numerous factors may affect the microbiological quality of pasteurized milk; these include duration of storage of raw milk prior to processing, heat-treatment and concentration of heat-resistant microorganisms, contaminants, packaging system adopted, post pasteurization storage conditions [21] and biofilm microflora; which seems to be the main recontamination source of milk [22]. Bacteria surviving pasteurization and cleaning in place (CIP) (circulation of cleaning liquids through machines and other equipment in a cleaning circuit) can potentially attach to piping and fitting surfaces, where they could promote the development of biofilm structures that enable protection against high temperatures and chemical compounds; as reported in two recent Algerian studies [23,24] ; showing that most bacterial types were found at different sampling points of pasteurization. Furthermore, biofilms are difficult to eradicate using conventional cleaning and disinfection procedures due to their resistant phenotype and disinfectants do not penetrate the biofilm matrix [21]. Coliform in milk is one of the best indications for judging its sanitary quality [20], thus when in high numbers, coliforms can cause food poisoning [14]. In our study, 63% of samples showed the presence of high level of total coliforms, in disagreement with the Algerian regulation, the rest of milk are considered satisfactory. In Tiaret

district, Aggad *et al.*[5] also found a high incidence (31%) of TC in pasteurized cow milk; as well, Shaltout *et al.*[2125] and Kunda *et al.*[2226] reported an elevated range (22 and 57%, in Jordan and Zambia, respectively). In contrast, Kunda *et al.*[2327] revealed a very low count of coliforms (4.8%) in raw milk, because of good hygienic practice. A possible explanation of the high level of TC is the milk contamination after processing; during packaging, or the use of poor quality packaging material. In addition, a probable contamination by polluted added water, after butter extraction, can occur. In fact, the presence of fecal coliforms reflects a recent fecal contamination, because these bacteria live primarily in the external environment [5,2428]. Nevertheless, according to Afif *et al.*[2529], the existence of coliforms may not necessarily indicate a direct fecal contamination of milk, but more precisely as an indicator of poor hygiene and sanitary practices during milking and further handling.

Fecal coliforms were present in 41% of total samples; while, Aggad *et al.*[5] by checking pasteurized milk in Tiaret district found only 2% of contamination. The presence of these pathogens with a high level seems to be linked to several factors; like failure of pasteurization, the personnel hygiene default, the failure of decontamination protocol of equipment and premises, and poor conditions of storage or milk protection. Furthermore, germs' ability to withstand pasteurization, the thermophilic bacteria can limit the shelf life of pasteurized milk [2630]. According to Guiraud [11] their presence at an abnormal rate indicates a poor general hygiene.

S. aureus was present in 61% of the checked milks; which is very considerable. Unlike, Leite *et al.*[2731] did not detect *S. aureus* in the pasteurized milk in Salvador. Nevertheless, Srairi *et al.*[2832] found 30% of contaminated raw milk, in Morocco. The presence of *S. aureus* in milk causes food poisoning; it is one of the most common causes of reported foodborne diseases, which is a major risk to public health [2933]. This sanitary problem can occur under certain conditions of heat-stable enterotoxin that can withstand heat treatments [3034] or when heat treatment process is ineffective; faulty pasteurization will not destroy all foodborne pathogens [31-3335-37]. Thus, according to Oliver *et al.*[3337] the increasing number of reports on detection of foodborne pathogens in pasteurized fluid milk and ready-to-eat dairy products clearly indicates that pasteurization alone is not the final solution for the control of milk-borne pathogens.

The SRC are reported in 34% checked samples. These germs are used as hygiene controls in the microbiological analysis of a number of food products. Their presence in pasteurized milk reflects a failure of pasteurization or contamination after pasteurization. Most of thermophilic bacteria grow slowly in refrigerated milk and are generally outgrown by Gram-negative psychrotrophic species that gain entry primarily as post-pasteurization contaminants [3438]. However, in the absence of psychrotrophic bacteria or if large numbers of thermophilic bacteria survive pasteurization; certain thermophilics particularly psychrotrophic spore forming *Bacillus spp.* can grow and cause milk spoilage [3539].

The fecal streptococci are potential fecal contamination agents; they are the most commonly used alternative or adjunct to coliform bacteria as faecal pollution indicators [3640]. We found these pathogens in all checked samples; which means that contamination occurred, on a large scale, before and / or during milk processing. Fecal streptococci may play a role in food poisoning [20].

At last, with regard to its poor hygienic quality, it appears clearly that checked pasteurized and marketed milk in Tiaret district is strongly similar to the raw milk collected in bad hygienic condition around world [25, 31,37,38,3929,35,41-43].

CONCLUSION

Pasteurization of milk assures safety for human consumption by reducing the number of live pathogenic bacteria present; thus, the public health benefits of pasteurization are well established. But, our study revealed that checked samples of pasteurized milks were not in compliance with sanitary regulation and its abundance with potential harmful germs constitute a serious health hazard. It is difficult to know if contamination has occurred after an effective pasteurization or it is due to failure of the process.

Consequently, urgent measures are required from the side of the responsible authorities to ensure pasteurization process effectiveness and to survey chain milk production.

Conflict of interests

None.

REFERENCES

1. Anonymous, 2010. Basic dairy bacteriology. Dairy Foods Science Notes. Department of Food Science Stocking Hall, Cornell University, Ithaca, NY, pp: 10.
2. MPI NZ (Ministry for Primary Industries, New Zealand), 2013. An Assessment of the Effects of Pasteurisation on Claimed Nutrition and Health Benefits of Raw Milk MPI Technical Paper No: 2014/13 ISBN No: 978-0-478-43209-1. ISSN New Zealand Government No: 2253-3923, October.
3. Fleming, D.W., S.L. Cochi, K.L. Mac Donald, J. Brondum, P.S. Hayes, B.D. Plikaytis and J.P. Giraud, 1998. Food microbiology, main food products microbiology. Ed., Dunod Paris, pp: 652.
4. Aggad, H., F. Mahouz, Y. Ahmed Ammar and M. Kihal, 2009. Evaluation of milk hygienic quality in Western Algeria. *Rev. Méd. Vét.*, 160 (12): 590-595.
5. Aggad, H., M. Bridja, AeK. Bouhai, M. Benaouali and A. Djebli, 2010. Some Quality Aspects of Pasteurized Milk in Algeria. *World Journal of Dairy and Food Sciences*, 5 (1): 21-24.
6. Labioui, H., L. Elmoualdi, A. Benzarouka, M. Elyachioui and M. Bernyouhssine, 2009. Etude physico-chimique et microbiologique des laits crus. Ed., bull. Soc. Pharm.Bordeaux, pp: 07-16.
7. Mathieu, J., 1998. Initiation de la physico-chimie du lait. Edition technique et documentation. Ed., Lavoisier, Paris, pp : 220.
8. Abiazar, R., 2007. Complication des protéines lactières par les extraits verts de caroubier. Propriétés technologiques des coagulums obtenus. Ed., Agroparistech Ecole Doctorale Abies, Paris, pp : 76.
9. JORA (Journal Officiel de la République Algérienne), 2013. Arrêté ministériel Du 16 aout 2012 rendant obligatoire la méthode de détermination de la teneur en matière sèche dans le lait, la crème et le lait concentré non sucré. N° 54, pp : 25-27.
10. Audigier, C.I., J.F. Figarella and C. Zonszain, 1980. Biochemical analysis engineering. 4 publishing. Publishing Doin Ed., Paris, pp. 265.
11. Giraud, J.P., 1998. Food microbiology, main food products microbiology. Ed., Dunod Paris, pp: 652.
12. JORA (Journal Officiel de la République Algérienne), 1998. Interministerial Decree. Specifications of some foodstuffs. N°35, pp. 199.
13. Catsaras, M. and C. Bourgeois, 1980. Les indices de contamination fécale : Techniques d'analyses et de contrôle dans les industries agro-alimentaires. Le contrôle microbiologique Vol 3. Tec et Doc Lavoisier. APRIA, pp : 331.
14. Joffin, C. and J.N. Joffin, 1999. Food microbiology. Collection biology and technique 5th publishing, pp: 211.
15. JORA (Journal Officiel de la République Algérienne), 2004. Ministerial Decree. Making obligatory a microbiological control method in pasteurized milk. N°70, pp: 19-22.
16. Schmidt, K.A., J. Stupar, J.E. Shirley, S. Adapa and D. Sukup, 1996. Factors affecting titratable acidity in raw milk Dairy Day, 1996, Kansas State University, Manhattan, KS, 60-62. URI: <http://hdl.handle.net/2097/8804>
17. Vignola, C.L., 2002. Science et technologie du lait: transformation du lait. Ed., presse international polytechnique, Ecole Polytechnique de Montréal, pp: 26.
18. Veisseyre, R., 1975. Dairy technology: Building, harvesting, treatment and transformation. Ed., La Maison Rustique Paris, pp: 230.
19. Bourgeois, C.M., J.F. Mescele and J. Zucca, 1996. Food Microbiology (Tome 01); microbiological aspect of safety and quality of food. Publishing technique and documentation Lavoisier Paris, pp: 272-292.
20. Guiraud, J.P. and JP Rosec, 2004. Pratique des normes en microbiologie alimentaire. Ed., AFNOR, pp: 298.
21. Sarkar, S., 2015. Microbiological Considerations: Pasteurized Milk. *International Journal of Dairy Science*, 10 (5): 206-218.
22. Marchand, S.,J. De Block, V. De Jonghe,A. Coorevits,M. Heyndrickx and L. Herman, 2012. Biofilm Formation in Milk Production and Processing Environments; Influence on Milk Quality and Safety. *Rev. Food. Sci. Food Safety*, 11: 133-147.
23. Malek, F., B. Moussa-Boudjemâa, F. Khaouani-Yousfi, A. Kalai and M. Kihel, 2012. Microflora of biofilm on Algerian dairy processing lines: An approach to improve microbial quality of pasteurized milk. *Afr. J. Microbiol. Res.*, 6(17): 3836-3844.

24. Cherif-Antar, A., B. Moussa-Boudjemâa, N. Didouh, K. Medjahdi, B. Mayo and A.B. Flórez, 2016. Diversity and biofilm-forming capability of bacteria recovered from stainless steel pipes of a milk-processing dairy plant. *Dairy Sci. & Technol.*, 96: 27–38.
25. Shaltout, F., H. Abdel-Samei, Y. Al-Tarazi and A. Al-Zamil, 2003. Sanitary status of raw cow milk marketed in northern Jordan. *Assiut Vet. Med. J.*, 49: 180-194.
26. Kunda, B., G.S. Pandey and J.B. Muma, 2015. Compositional and sanitary quality of raw milk produced by smallholder dairy farmers in Lusaka Province of Zambia. *Livestock Res. Rural Dev.*, 27: 201.
27. Kunda, B., G.S. Pandey, C. Mubita, J.B. Muma and C. Mumba, 2015. Compositional and microbial quality of heat-treated milk brands marketed in Lusaka, Zambia. *Livestock Res. Rural Dev.*, 27: 143.
28. Douglas, J.R., 2003. Bulk tank cultures are the dairy man best friend. University of Wisconsin milking research and instruction lab (Report) N°2223 (Report N°2223).
29. Afif, A., M. Faid and M. Najimi, 2007. Effects of breeding and hygienic practices on raw cow milk quality in Tadla area, Morocco. *Livestock Res. Rural Dev.*, 19: 181.
30. TeGiffel, M.T., 1997. Isolation, identification and characterization of *Bacillus cereus* from the dairy environment. Thesis, Land Bouw Universiteit, Wageningen, ISBN: 90-5485- 694-7.
31. Leite, C.C., A.G. Guimarães, P.N. Assis, M.D. Silva and C.S. Andrade, 2002. Qualidade bacteriológica do leite integral (tipo C) comercializado em Salvador-Bahia. *Rev. Bras. Saúde Prod. Anim.*, 3: 21-25.
32. Sraïri, M.T., J. Moudnib, L. Rahho and A. Hamama, 2006. How do milking conditions affect the hygienic quality of raw milk? Case study from Moroccan dairy farms. *Livestock Res. Rural Dev.*, 18: 97.
33. Kadariya, J., T.C. Smith and D. Thapaliya, 2014. *Staphylococcus aureus* and Staphylococcal Food-Borne Disease: An Ongoing Challenge in Public Health. Art ID 827965, 9 p. <http://dx.doi.org/10.1155/2014/827965>
34. Larpent, J.P., 1997. *Microbiologie alimentaire : Technique de laboratoire*, Ed., Doc Tec, pp : 1072.
35. Jorgensen, H.J., T. Mork, H.R. Hogasen and L.M. Rorvik, 2005. Enterotoxigenic *Staphylococcus aureus* in bulk milk in Norway. *J. Appl. Microbiol.*, 99: 158-166.
36. Fagundes, H., L. Barchesi, A. Nader Filho, L. Menezes Ferreira, C. Augusto and F. Oliveira, 2010. Occurrence of *staphylococcus aureus* in raw milk produced in dairy farms in São Paulo State, Brazil. *Braz. J. Microbiol.*, 41: 376-380.
37. Oliver, S.P., B.M. Jayarao and R. Almeida, 2005. Foodborne Pathogens in Milk and the Dairy Farm Environment: Food Safety and Public Health Implications, a Review. *Foodborne Pathog. Dis.*, 2: 2. doi:10.1089/fpd.2005.2.115.
38. Cousin, M.A., 1982. Presence and activity of psychrotrophic microorganisms in milk and dairy products. *J. Food prot.*, 45: 172.
39. Overcast, W.W. and K. Atmaram, 1974. The role of *bacillus cereus* in sweet curdling of fluid milk. *J. Milk Food Technol.*, 37: 233.
40. Sinton, L.W., A.M. Donnison and C.M. Hastie, 1993. Fecal streptococci as fecal pollution indicators: A review. Part I: Taxonomy and enumeration, *New Zeal. J. Mar. Fresh.*, 27(1): 101-115.
41. Chye, F. Y., A. Abdullah and M K Ayob, 2004. Bacteriological quality and safety of raw milk in Malaysia. *Food Microbiol.*, 21(5): 535-541.
42. Schooman, L. and E.S. Swai, 2011. Marketing, handling and physical quality of raw marketed milk in Tanga region of Tanzania. *Livestock Res. Rural Dev.*, 23: 191.
43. Hidalgo-Milpa, M., E. Sánchez-Vera and A. Espinoza-Ortega, 2015. Quality of milk in a traditional milk-cheese chain in the highlands of central Mexico. *Livestock Res. Rural Dev.*, 27: 232.